

ECLIPSE®

Pour-On for Cattle



ACTIVES	RAINFAST	DOSE RATE	ORAL DRENCH	CONTROLS	WHP (days)				
					<table border="1"><tr><td>MEAT</td><td>MILK</td></tr><tr><td>35</td><td>35</td></tr></table>	MEAT	MILK	35	35
MEAT	MILK								
35	35								

ECLIPSE® Pour-On for Cattle is the world's first dual combination cattle pour-on which treats and controls internal and external parasites.

CONTAINS

10 g/L abamectin, 200 g/L levamisole.

Abamectin is a member of the macrocyclic lactones. It is effective against internal and external parasites sensitive to this family. Levamisole is a member of the imidazothiazole class of anthelmintics with efficacy against internal parasites sensitive to this family, including lungworm.

INDICATIONS

INTERNAL PARASITES

The product will treat and control sensitive strains of *Ostertagia ostertagi*, *O. lyrata*, *Cooperia oncophora*, *C. surnabad*, *C. punctata*, *Trichostrongylus axei*, mature *Trichuris* and adult and immature lungworm (*Dictyocaulus bovis*). Is effective against inhibited 4th stage larvae (L4) of *Ostertagia* and *Cooperia* species that can hide in the gut as well.

EXTERNAL PARASITES

The product will treat and control sensitive strains of both sucking (*Linognathus vituli*) and biting (*Damalinia bovis*) lice.

RECOMMENDED USE

Timing of treatments will depend on the parasite pressure on individual farms. Diagnostic testing and historical results including faecal egg counts (FECs), faecal egg count reduction tests (FECRTs) and larval cultures can help you to understand the parasite burden and resistance challenges you are facing.

Talk to your local vet about parasite management practices for your area to reduce the development of resistance. It is advisable that a resistance test be conducted regularly when using any parasite treatment.



PRECAUTIONS & NOTES

Do not use on animals which are severely stressed or in ill health. Care should be taken in handling pregnant animals. Do not exceed recommended dose rates. Doses of 3 or more times the recommended can cause symptoms of levamisole toxicity, so estimate liveweights carefully. Dehydrated animals may be more susceptible to toxicity. Fatal interactions occur between levamisole and organophosphate dips.

Do not use if it's raining or looks likely to rain within 2 hours of treatment.

Temporary hair loss occasionally occurs following application. Therefore, do not treat cattle prior to showing.

ECLIPSE®

Pour-On for Cattle

DOSAGE GUIDE – CATTLE

Dose rate is 1 mL/20kg body weight (0.5 mg/kg abamectin and 10 mg/kg levamisole). Apply along the backline from between the shoulders to the tail using the applicator. Check the accuracy of drench guns regularly.

BODY WEIGHT KG	DOSE mL	2.5 L TREATS	5 L TREATS	5.5 L TREATS
61 – 80	4	625	1250	1375
81 – 100	5	500	1000	1100
101 – 120	6	416	833	916
121 – 140	7	357	714	785
141 – 160	8	312	625	687
161 – 180	9	277	555	611
181 – 200	10	250	500	550
201 – 220	11	227	454	500
221 – 240	12	208	416	458
241 – 260	13	192	384	423
261 – 280	14	178	357	392
281 – 300	15	166	333	366
301 – 320	16	156	312	343
321 – 340	17	147	294	323
341 – 360	18	138	277	305
361 – 380	19	131	263	289
381 – 400	20	125	250	275
401 – 420	21	119	238	261
421 – 440	22	113	227	250
441 – 460	23	108	217	239
461 – 480	24	104	208	229
481 – 500	25	100	200	220
501 – 520	26	96	192	211
521 – 540	27	92	185	203
541 – 560	28	89	178	196
561 – 580	29	86	172	189
581 – 600	30	83	166	183
601 – 620	31	80	161	177
621 – 640	32	78	156	171
641 – 660	33	75	151	166

For cattle over 660 kg: Dose at 1 mL per 20 kg b.w.
(2.5 mL/50 kg bodyweight)

A representative sample of animals should be weighed before treatment either with scales or a weighband. Dose rate to be based on the heaviest cattle in each group. DO NOT UNDERDOSE. If there is a large variation in size within the group, draft into two or more lines based on bodyweight to avoid excessive overdosing.

PACK SIZE

Available in 2.5 L, 5 L, 5.5 L packs.
(applicator included)



PROUDLY AVAILABLE FROM YOUR LOCAL PARTICIPATING VETERINARY CLINIC

See product label for full claim details and directions for use. ECLIPSE® is a registered trademark of Boehringer Ingelheim Vetmedica GmbH, used under license. ACVM Registration No. A009270. © Copyright 2024 Boehringer Ingelheim Animal Health New Zealand Limited. All rights reserved. NZ-BOV-0006-2023.

